

Tests of possible non standard properties of the heavy quarks and the Higgs boson ¹

G.J. Gounaris^a, J. Layssac^b, D.T. Papadamou^a, G. Tsirigoti^a
and F.M. Renard^b

^aDepartment of Theoretical Physics, University of Thessaloniki,
Gr-54006, Thessaloniki, Greece.

^bPhysique Mathématique et Théorique, UPRES-A 5032
Université Montpellier II, F-34095 Montpellier Cedex 5.

Abstract

We discuss the sensitivity of the processes $e^+e^- \rightarrow b\bar{b}$, $e^+e^- \rightarrow t\bar{t}$ and $\gamma\gamma \rightarrow H$ to certain residual New Physics interactions affecting the heavy quark and Higgs boson sector. With a linear collider of 500 GeV , one should be able to detect and identify such effects with a characteristic scale up to 50 TeV .

¹Contribution to the studies on Physics and Detectors for the Linear Collider”, to appear in DESY 97-123E. Partially supported by the EC contract CHRX-CT94-0579.

1 Introduction

We assume that a certain dynamics generically called New Physics (NP) exists beyond the Standard Model (SM) and is responsible for many of its unexplained features like the structure of the scalar sector and the mass spectrum of the leptons, the quarks and the gauge bosons. It is further assumed that this new dynamics involves heavy degrees of freedom characterized by an effective scale much higher than the electroweak scale; i.e. $\Lambda \gg M_W$. It is then natural to expect that this dynamics which leads to the mass of the usual particles, also generates some additional new interactions among these particles, called residual NP effects. This picture suggests that the NP effects should most probably affect particles especially concerned by the mass generation mechanism, namely the heavy quarks and the W_L , Z_L and scalar states. Inevitably, such NP will also involve "anomalous" gauge interactions, which are required by the gauge principle whenever derivative interactions appear.

Recent hints for anomalous effects have appeared in $Z \rightarrow b\bar{b}$; i.e. $\Gamma(Z \rightarrow b\bar{b})$ is 1.8σ too high and A_b is 2σ too low, as compared to SM predictions, [1]. There also exist unexplained features in the high p_T jets produced at Tevatron [2] and in high q^2 events at HERA [3].

To describe these residual effects in a most model-independent way, we use the effective lagrangian method. Thus, for $M_W \lesssim E \ll \Lambda$, by integrating out all NP degrees of freedom, one obtains an effective Lagrangian describing the residual effects among usual particles. This Lagrangian satisfies of course $SU(3) \times SU(2) \times U(1)$ gauge invariance broken at the electroweak scale, and involves a Higgs particle which should also be of the order of the electroweak scale or heavier. For sufficiently large Λ , the lowest dimension operators ($d = 6$) should dominate NP contribution. One is then able to draw a list of operators [4], $\mathcal{O}_i$, being associated to corresponding dimension ($4 - d = -2$) couplings f_i . These couplings are the free parameters of the description and, through unitarity considerations, they can be related to the NP scale where either new strong interactions are generated or new particles are produced [5, 6].

The general description of NP in terms of the list $\mathcal{O}_i$ involves various classes of bosonic and fermionic operators with specific features. Bosonic operators are usually separated into "non-blind", "blind" and even "super-blind" ones, depending on their appearance in Z-peak observables at tree or loop levels, [7, 8, 9]. The same concept has been extended to the list of heavy quark operators; i.e. operators involving quarks of the third family. In classifying them, the possible presence of the t_R field has been used [10, 11]. To reduce the number of the independent operators, the equations of motion for 3rd-family quarks and scalar fields have been used, since they do not mix other families, to the extent that all fermion masses except the top are neglected. Thus finally we end up with three classes of such operators involving quarks of the third family [12].

Tests of purely bosonic operators in e^+e^- annihilation processes were discussed in previous reports, (for a review see [13, 14]). In Section 2 we report on the tests concerning the heavy quark sector, and Section 3 is devoted to an example of tests concerning operators inducing CP violating $H\gamma\gamma$ interactions.

2 Tests in the heavy quark sector

2.1 Tests in $e^+e^- \rightarrow t\bar{t}$

Obviously, the heavy top should be a privileged place for looking for these effects; so we start with the process $e^+e^- \rightarrow t\bar{t}$ observable at LC [14]. The aforementioned operators lead to modifications of the $\gamma t\bar{t}$ and $Z t\bar{t}$ couplings. The general form of such CP-conserving vertices is given by [15, 16, 17, 11]

$$-i\epsilon_\mu^V J_V^\mu = -ie_V \epsilon_\mu^V \bar{u}_t(p) [\gamma^\mu d_1^V(q^2) + \gamma^\mu \gamma^5 d_2^V(q^2) + (p-p')^\mu d_3^V(q^2)/m_t] v_{\bar{t}}(p') , \quad (1)$$

where ϵ_μ^V is the polarization of the vector boson $V = \gamma, Z$. The outgoing momenta (p, p') refer to $(t, \bar{t})$ respectively and satisfy $q \equiv p + p'$. The normalizations are determined by $e_\gamma \equiv e$ and $e_Z \equiv e/(2s_W c_W)$, while d_i^V are in general q^2 dependent form factors. At tree level SM feeds the first set, called set (1), of the three couplings

$$d_1^{\gamma, SM0} = \frac{2}{3} , \quad d_1^{Z, SM0} = g_{Vt} = \frac{1}{2} - \frac{4}{3}s_W^2 , \quad d_2^{Z, SM0} = -g_{At} = -\frac{1}{2} . \quad (2)$$

SM at 1-loop and NP lead to additional (q^2 -dependent) contributions to these couplings, as well to the couplings of the second set, called set (2), consisting of $d_2^V(q^2)$, $d_3^V(q^2)$ and $d_3^Z(q^2)$.

Departures from the SM (tree + 1-loop) are defined as:

$$\bar{d}_j^V \equiv d_j^V - d_j^{V, SM} . \quad (3)$$

The explicit expressions of the contributions to the $\bar{d}_j^V$ for each $\mathcal{O}_i$ at tree level and at 1-loop are given in [11]. We will now precisely determine the accuracy at which they can be observed or constrained.

The reaction $e^+e^- \rightarrow t\bar{t}$ offers a way to determine all six couplings by studying the top quark spin density matrix elements. There are six independent such elements $\rho_{++,--,+-}^{L,R}$, (where L, R refer to the longitudinal polarization of the e^- beam), which can be reconstructed through the decay chains $t \rightarrow Wb \rightarrow l\nu b$ and $t \rightarrow Wb \rightarrow q\bar{q}'b$, for any q^2 and production angle θ [11, 17]. Measurements of the magnitude of density matrix elements (like the production cross section $\sigma_{t\bar{t}}$) are affected by the lack of precise knowledge of the $t \rightarrow Wb$ decay width [18, 19]. More accurate are the measurements of asymmetries like forward-backward or L-R asymmetries, which, being ratios of cross sections, are free of this uncertainty. In order to study the $\gamma t\bar{t}$ and $Z t\bar{t}$ couplings, we have tried to rely as much as possible on such observables.

In the general 6-parameter case this is not fully possible. Asymmetries only depend on two combinations of the first set of the three $\bar{d}_j^V$ couplings. To get a third one, information sensitive to the absolute normalization, like $\sigma_{t\bar{t}}$, is required (which also depends on the top quark decay width). For the second set of couplings we do not have this problem, since asymmetries are now sensitive to ratios of the type $\bar{d}/d^{SM}$ and allow to constrain the complete set (2). We have made applications to an LC collider [14] with 0.5, 1 and

2 TeV and luminosity of 20, 80 and 320 fb^{-1} respectively, leading to more than 10^4 events. A reconstruction and detection efficiency of 18% [19] has been applied before computing the statistical accuracy for each observable. From this we have obtained the accuracy expected in the determination of the top quark density matrix elements and the constraints on the NP couplings $\bar{d}_j$ (ellipsoid in 6-parameter space). Examples are shown in Fig.1a,b. The complete set of results is given in [17].

The essential features are the following. For couplings of set (1), a band of width ± 0.02 appears, as information on the overall magnitude of the amplitudes is missing when only asymmetries are used. When cross section measurements are added assuming an uncertainty of 20% (2%) on the top decay width, a constraint of the order of ± 0.05 (± 0.02) arises. On the contrary, for couplings of set (2), asymmetries alone constrain the couplings at the level of ± 0.02 in the unpolarized case and at the level of ± 0.01 in the polarized one.

We have also studied constrained cases like the 4-parameter one obtained by restricting to the NP forms generated by the operators $\mathcal{O}_{t2}$, $\mathcal{O}_{Dt}$, $\mathcal{O}_{tW\Phi}$ and $\mathcal{O}_{tB\Phi}$ contributing to the $\gamma t\bar{t}$ and $Z t\bar{t}$ vertices at tree level [11]. This case is further reduced to a three parameter one if only the operators $\mathcal{O}_{t2}$, $\mathcal{O}_{tW\Phi}$ and $\mathcal{O}_{tB\Phi}$ generated in the dynamical models studied in [12] are retained. Finally, a two parameter case is met if one insists on a chiral form for NP where only anomalous left-handed and right-handed $Z t\bar{t}$ couplings appear [20].

In these constrained cases it is sufficient to use only asymmetries, and the typical accuracy for both sets of couplings is of the order of ± 0.02 in the no polarization case, and of the order of ± 0.01 when polarization is available. Finally, the results for all operators listed in Class 1 [12], can be found in Table 7 of [17]. Essentially there are two levels of constraints. For the four operators $\mathcal{O}_{t2}$, $\mathcal{O}_{Dt}$, $\mathcal{O}_{tW\Phi}$ and $\mathcal{O}_{tB\Phi}$ contributing at tree level at the LC, the constraints reach effective scales lying in the 5 to 50 TeV range, which is much larger than the range reached through the indirect result arising from the Z-peak contributions. On the contrary for the operators $\mathcal{O}_{qt}$, $\mathcal{O}_{qt}^{(8)}$, $\mathcal{O}_{tb}$ contributing at loop level to $e^-e^+ \rightarrow t\bar{t}$, the Z-peak constraints are not expected to be substantially improved, and the NP scales should lie in the 1-5 TeV range. However for the two operators $\mathcal{O}_{tt}$ and $\mathcal{O}_{tG\Phi}$, which contribute only at loop level and do not give substantial effects at Z-peak, interesting constraints in the 5-10 TeV range will arise from $e^-e^+ \rightarrow t\bar{t}$.

2.2 Tests in $e^+e^- \rightarrow b\bar{b}$

We now discuss the $e^+e^- \rightarrow b\bar{b}$ channel. For the operators in Class 1 and Class 2 [12], the aforementioned channel at LC cannot improve the existing information by other means. Thus for the Class 2 operators which give tree level contributions to the $\gamma b\bar{b}$ and $Z b\bar{b}$ couplings [21], the 0.5% accuracy obtained from the $Z \rightarrow b\bar{b}$ measurements pushes the NP scale in the 10 TeV range; and this cannot be improved by new measurements in the $e^+e^- \rightarrow b\bar{b}$ channel at higher energies. This is also true for the operators of Class 1, which contribute only at loop level to the $\gamma b\bar{b}$ and $Z b\bar{b}$ vertices. Some of these operators are in fact constrained by Z-peak measurements, because their contribution there is enhanced by m_t^2/M_W^2 factors [10]. We have also seen that in some cases, like for $\mathcal{O}_{qt}$, $\mathcal{O}_{qt}^{(8)}$ and $\mathcal{O}_{tb}$,

Z peak measurements give better constraints than $e^+e^- \rightarrow t\bar{t}$ at high energies.

Nevertheless, we have found that $e^+e^- \rightarrow b\bar{b}$ measurements beyond Z-peak can improve the constraints for certain "non-blind" Class 3 operators containing derivatives of gauge fields [12]. This turns out to be the case for the gauge boson operators $\mathcal{O}_{DW}$, $\mathcal{O}_{DB}$ studied in [22, 23], and the operators $\mathcal{O}_{qW}$, $\mathcal{O}_{qB}$, $\mathcal{O}_{bB}$ studied in [24]. They all give effective four-fermion $e^+e^-b\bar{b}$ contact interactions, whose contribution is enhanced with respect to the Z, γ ones, by q^2 factors. This property allows the disentangling of these contributions from those of all other operators, by using the so-called "Z-peak subtraction method" [25]. This method consists in using as inputs $\Gamma(Z \rightarrow f\bar{f})$, A_f instead of G_μ , s_W^2 . Thus, it automatically absorbs the NP effects at Z peak and leaves only room for those ones which can survive in the difference of structure functions of the type

$$F_i(q^2) - F_i(M_Z^2) \quad , \quad (4)$$

which are combinations of self-energies, vertices, box and NP contributions. This way, only the contributions from the aforementioned Class 3 operators $\overline{\mathcal{O}}_{DB}$, $\overline{\mathcal{O}}_{DW}$, $\mathcal{O}_{qW}$, $\mathcal{O}_{qB}$, $\mathcal{O}_{bB}$ survive sufficiently beyond the Z-peak. The operators $\overline{\mathcal{O}}_{DB}$, $\overline{\mathcal{O}}_{DW}$ are constrained from $e^+e^- \rightarrow l^+l^-$ and $e^+e^- \rightarrow q\bar{q}$ and have been studied in [22, 23]; while $\mathcal{O}_{qW}$, $\mathcal{O}_{qB}$, $\mathcal{O}_{bB}$ can be studied by measuring the observables σ_b , A_{FB}^b , A_{LR}^b , $A_{FB}^{pol(b)}$, which of course require polarized beams [24]. At LEP2 polarization is not available and we only get two constraints for these three operators. This is the origin of the band appearing in Fig.2; (illustrations for other couples of operators can be found in ref.[24]). For polarized beams at LC, the system can be completely constrained leading to sensitivity limits on NP scales in the range 30-50 TeV; indeed a rather high level.

In conclusion, we summarize the constraints obtained or expected from for the whole set of $dim = 6$ operators. We give below the range of NP scales (in TeV) up to which experiments at LEP and LC can be sensitive.

Bosonic operators		
non-blind	17 (LEP2)	50 (LC)
blind (TGC)	1.5 (LEP2)	20 (LC)
superblind(Higgs)	7 (LEP2)	20 (LC), 70 (LC $\gamma\gamma$)

Heavy quark operators		
Class 1	5 (LEP1)	10-50 (LC)
Class 2	10 (LEP1)	10 (LC)
Class 3	8 (LEP2)	30-50 (LC)

As one can see, limits expected from the LC should improve by one order of magnitude the ones indirectly obtained at LEP1 or expected from LEP2. They should lie in the several tens of TeV range. This is a domain which covers various types of models beyond SM (Technicolour, extended gauges,...). Our hope is that experiments in various sectors will reveal certain correlations which could select a few operators of our list and give hints for the structure of new physics.

3 Tests of CP violating $H\gamma\gamma$ interactions

The possibility to have polarized e^- , e^+ beams in a linear collider, renders them very efficient in searching for CP violation among the NP interactions. For example, by measuring $e^-e^+ \rightarrow W^-W^+$ using polarized beams, it is possible to improve considerably [26], the present constraints on the CP violating triple gauge boson interactions [27]. Moreover at a LC, it is possible to search for CP violating forces directly connected to the Higgs particle [28, 29, 30].

Here we concentrate on the option to use LC as a $\gamma\gamma$ collider by backscattering polarized laser photons from polarized electrons and positrons [31]. In particular we concentrate on studying the CP violating forces affecting the $\gamma\gamma \rightarrow H$ process. In the philosophy followed in the present note, where all new physical degrees of freedom are too heavy to be directly produced, such CP violation may arise from the $dim = 6$ operators called $\tilde{\mathcal{O}}_{WW}$, $\tilde{\mathcal{O}}_{BB}$ in [28, 32, 33], whose couplings we denote by $\bar{d}$ and $\bar{d}_B$ respectively.

As soon as the Higgs particle is discovered, a general study of all Higgs properties, including of course the possibility for CP violating forces, will become mandatory. For CP violation in particular, such studies are based on constructing appropriate spin asymmetries with respect to the polarizations of the laser photons and e^- , e^+ beams. For Higgs production through $\gamma\gamma$ collisions these are constructed by noting that for a laser photon described by the density matrix

$$\rho_{laser}^N = \frac{1}{2} \begin{pmatrix} 1 + P_\gamma & -P_t e^{-2i\varphi} \\ -P_t e^{+2i\varphi} & 1 - P_\gamma \end{pmatrix} \quad (5)$$

in its helicity basis, the corresponding photon obtained by backscattering the laser one from the e^- beam is described by the density matrix [31]

$$\rho^{BN} = \frac{1}{2} \begin{pmatrix} 1 + \xi_2(x) & -\xi_{13}(x) e^{-2i\varphi} \\ -\xi_{13}(x) e^{+2i\varphi} & 1 - \xi_2(x) \end{pmatrix} , \quad (6)$$

where P_γ describe the circular polarization of the laser photon, (P_t, φ) give the magnitude of the linear polarization and its azimuthal angle around the momentum of laser-photon, while the Stokes parameters ξ_j of the backscattered photon are known functions of P_γ , P_t , φ and of the (longitudinal) polarization P_e of the e^- beam [31]. Corresponding quantities (denoted as barred below) are analogously defined for the photon backscattered from the positron also.

The two possible asymmetries are ($\chi \equiv \varphi - \bar{\varphi}$)

$$\tilde{A}_{lin} = \frac{|N_{\tau_H}(\chi = \frac{\pi}{4}) - N_{\tau_H}(\chi = -\frac{\pi}{4})|}{N_{\tau_H}(\chi = \frac{\pi}{4}) + N_{\tau_H}(\chi = -\frac{\pi}{4})} = \frac{\langle \xi_{13} \bar{\xi}_{13} \rangle}{1 + \langle \xi_2 \bar{\xi}_2 \rangle} A_{lin} \quad (7)$$

which uses the relative angles of the linear polarizations of the two photons, and

$$\tilde{A}_{circ} = \frac{|N_{\tau_H}^{++} - N_{\tau_H}^{--}|}{N_{\tau_H}^{++} + N_{\tau_H}^{--}} = \frac{|\langle \xi_2 + \bar{\xi}_2 \rangle|}{1 + \langle \xi_2 \bar{\xi}_2 \rangle} A_{circ} , \quad (8)$$

where the signs in the upper indices describe generically two opposite signs for the pairs (P_e, P_γ) and $(\bar{P}_e, \bar{P}_\gamma)$, which generates an asymmetry between two opposite circular polarizations for the photons backscattered from e^- and e^+ respectively.

The quantities A_{lin} and A_{circ} defined in (7,8), are sensitive to the combination $\bar{d}s_W^2 + \bar{d}_b c_W^2$ of the NP couplings, and depend of course also on the SM dynamics. It turns out that A_{circ} is useful only for $200 GeV \lesssim m_H$; while A_{lin} can be useful for both $100 GeV \lesssim m_H \lesssim 150 GeV$ as well as for $200 GeV \lesssim m_H$ [33]. The reachable sensitivity limits are indicated in Tables I, II (where $x_0 \equiv 4E\omega_0/m_e^2 \leq 2(1 + \sqrt{2})$ is the usual parameter determined from the laser energy) and background effects have to some extent been taken into account [33].

It is concluded from this that using polarized beams for realizing the $\gamma\gamma$ colliders, it is possible to construct observables sensitive only to the CP violating NP couplings and thereby distinguish them from the CP conserving ones. This is not attainable for unpolarized beams. Thus, the process $\gamma\gamma \rightarrow H$ at a $0.5 TeV$ tunable linear collider may be able to put limits on the NP coupling $\bar{d}s_W^2 + \bar{d}_B c_W^2$ at the $10^{-3} - 10^{-4}$ level, for Higgs masses in the ranges $100 \lesssim m_H \lesssim 150 GeV$ and $200 \lesssim m_H \lesssim 350 GeV$. This way, one can probe NP scales in the range of 10-20TeV for NP generated by $\tilde{\mathcal{O}}_{WW}$, and in the range of 30-50TeV in the case $\tilde{\mathcal{O}}_{BB}$.

This information is complementary, and at least an order of magnitude more precise than the one attainable through production of WW pairs in $\gamma\gamma$ collisions, where the same combination of NP couplings is measured [32, 34, 30]. Information on independent combinations of the CP violating couplings at the level of 10^{-2} may be obtained by looking at $e^-e^+ \rightarrow H\gamma, HZ$ [6, 28, 29]. Thus, a combination of such measurements should be able to constrain separately each of the two CP violating couplings $\bar{d}$ and $\bar{d}_B$ at the 10^{-2} level [33].

References

- [1] See the talks of A. Blondel and M.W. Grünewald at the Warsaw conference on High Energy Physics (1996).
- [2] CDF, F.Abe et al, Phys. Rev. Lett. **77** (1996) 438; Phys. Rev. **D55** (1997) 5263.
- [3] H1 Coll., C. Adloff et al, Z. f. Phys. **C74** (1977) 191; ZEUS Coll, J. Breitweg et al, Z. f. Phys. **C74** (1997) 207.
- [4] W. Buchmüller and D. Wyler, Nucl. Phys. **B268** (1986) 621; C.J.C. Burgess and H.J. Schnitzer, Nucl. Phys. **B228** (1983) 454; C.N. Leung, S.T. Love and S. Rao Z. f. Phys. **C31** (1986) 433. C. Arzt, M.B. Einhorn and J. Wudka, Nucl. Phys. **B433** (1995) 41.
- [5] G.J. Gounaris, J. Layssac and F.M. Renard, Phys. Lett. **B332** (1994) 146; G.J. Gounaris, J. Layssac, J.E. Paschalis and F.M. Renard, Z. f. Phys. **C66** (1995) 619; M. Hosch, K. Whisnant and B-L. Young, AMES-HET-96-04 (1996).

TABLE I. 3σ upper bounds on CP-violating NP couplings from A_{lin} asymmetry, using $H \rightarrow b\bar{b}$ for the two polarization choices:
 $P_t = \bar{P}_t = 1, P_\gamma = \bar{P}_\gamma = 0, \quad (P_t = \bar{P}_t = P_\gamma = \bar{P}_\gamma = 1/\sqrt{2}, P_e = \bar{P}_e = 1).$

m_H (GeV)	x_0	upper limit on $\bar{d}_B c_W^2 + \bar{d}_S^2$	$(\varphi - \bar{\varphi})$ -averaged Events
100	0.5	10^{-3} (2×10^{-3})	197 (168)
	0.8	2×10^{-2} (1.6×10^{-2})	184 (233)
	1	0.12 (0.05)	160 (255)
120	0.5	5×10^{-4} (8.5×10^{-4})	200 (160)
	0.8	8×10^{-3} (6×10^{-3})	207 (223)
	1	0.05 (0.017)	192 (264)
	1.5	0.16 (0.11)	175 (306)
140	0.5	4×10^{-4} (6.5×10^{-4})	144 (111)
	0.8	5×10^{-3} (3.2×10^{-3})	159 (143)
	1	0.03 (0.01)	156 (181)
	1.5	0.09 (0.06)	147 (237)
	2	0.22 (0.22)	138 (252)
150	0.5	5×10^{-4} (8×10^{-4})	85 (64)
	0.8	4.3×10^{-3} (3×10^{-3})	106 (87)
	1	0.023 (0.008)	107 (113)
	1.5	0.08 (0.05)	102 (157)
	2	0.2 (0.2)	97 (174)

TABLE II. 3σ upper bounds on CP-violating NP couplings from A_{circ} asymmetry, using $H \rightarrow ZZ \rightarrow l^- l^+ X$ decay and circularly polarized laser beams with $P_e = \bar{P}_e = -P_\gamma = -\bar{P}_\gamma = \pm 1$.

m_H (GeV)	x_0	upper limit on $\bar{d}_B c_W^2 + \bar{d}_S s_W^2$	ξ_2 -averaged Events
200	1	5.7×10^{-4}	141
	1.5	6×10^{-4}	133
	2	6.2×10^{-4}	121
	2.5	6.5×10^{-4}	112
	4	7×10^{-4}	95
	4.82	7.3×10^{-4}	89
240	1	6.3×10^{-4}	49
	1.5	4.2×10^{-4}	109
	2	4.3×10^{-4}	105
	2.5	4.4×10^{-4}	99
	4	4.7×10^{-4}	85
	4.82	5×10^{-4}	80
280	1.5	4.5×10^{-4}	55
	2	3.7×10^{-4}	78
	2.5	3.7×10^{-4}	78
	4	4×10^{-4}	70
	4.82	4×10^{-4}	66
320	2	4.8×10^{-4}	31
	2.5	3.6×10^{-4}	53
	4	3.5×10^{-4}	56
	4.82	3.6×10^{-4}	54
350	2.5	6×10^{-4}	15
	4	3.4×10^{-4}	45
	4.82	3.4×10^{-4}	46

- [6] G.J. Gounaris, F.M. Renard and N.D. Vlachos, Nucl. Phys. Nucl. Phys. **B459** (1996) 51.
- [7] A. De Rújula *et.al.* , Nucl. Phys. **B384** (1992) 3.
- [8] K.Hagiwara et al, Phys. Lett. **B283** (1992) 353, Phys. Rev. **D48** (1993) 2182; K. Hagiwara, S. Ishihara, R. Szalapski and D. Zeppenfeld, Phys. Lett. **B283** (1992) 353, Phys. Rev. **D48** (1993) 2182.
- [9] G.J. Gounaris, F.M. Renard and G. Tsirigoti, Phys. Lett. **B338** (1994) 51, Phys. Lett. **B350** (1995) 212.
- [10] G. Gounaris, F.M. Renard and C. Verzegnassi, Phys. Rev. **D52** (1995) 451.
- [11] G.J. Gounaris, M. Kuroda and F.M. Renard, PM/96-22, THES-TP 96/06, Phys. Rev. **D54** (1996) 6861.
- [12] G.J. Gounaris, D. Papadamou and F.M. Renard, PM/96-28, THES-TP 96/09, hep-ph/9609437, to appear in Z.f. Physik C; G.J. Gounaris, D.T. Papadamou and F.M. Renard, PM/96-31, THES-TP 96/10, hep-ph/9611224, (unpublished).
- [13] Physics at LEP2, Proceedings of the Workshop-Geneva, Switzerland (1996), CERN 96-01, G. Altarelli, T. Sjostrand and F. Zwirner eds.
- [14] e^+e^- Collisions at 500 GeV: The Physics Potential, Proceedings of the Workshop-Munich, Annecy, Hamburg, DESY 92-123A (1992), 92-123B (1992), 93-123C (1993), P.M. Zerwas ed.
- [15] K. Whisnant, J.M. Yang, B.-L. Young and X. Zhang, AMES-HET-97-1, hep-ph/9702305; A. Data and X. Zhang, hep-ph/9611247.
- [16] C.-P. Yuan, Lectures at the VI Mexican School of Particles and Fields, Villahermosa, Mexico 1994, MSUHEP-50228.
- [17] G.J. Gounaris, J. Layssac and F.M. Renard, Phys. Rev. **D55** (1997) 5786.
- [18] For a review, see e.g., J.H. Kühn, TTP96-18 (1996); F. Larios, E. Malkawi and C.-P. Yuan, MSUHEP-60922, hep-ph/9609482; D.O. Carlson and C.-P. Yuan, MSUHEP-50823, 1995; C.-P. Yuan hep-ph/9604434.
- [19] R. Frey, talk given at Conf. on Phys. and Exp. with Linear Colliders, Morioka-Appi, Japan (1995), OREXP 96-04.
- [20] R.D. Peccei and X. Zhang, Nucl. Phys. **B337** (1990) 269; R.D. Peccei, S. Peris and X. Zhang, Nucl. Nucl. Phys. **B349** (1991) 305; E. Malkawi and C.-P. Yuan, Phys. Rev. **D52** (1995) 472; F. Larios and C.-P. Yuan, hep-ph/9606397.

- [21] X.Zhang and B.L. Young, Phys. Rev. **D51** (1995) 6584; A. Datta, K. Whishnant, Bing-Lin Young and X. Zhang, preprint AMES-HET-97-6.
- [22] K. Hagiwara, S. Matsumoto, and R. Szalapski, Phys. Lett. **B357** (1995) 411; K. Hagiwara, T. Hatsukano, S. Ishihara and R. Szalapski, KEK-TH-497(1996).
- [23] A. Blondel, F.M. Renard, L. Trentadue and C. Verzegnassi, Phys. Rev. **D54** (1996) 5567; F.M. Renard and C. Verzegnassi, PM/96-27(1996), to appear in Phys.Rev.D.
- [24] G.J. Gounaris, D.T. Papadamou and F.M. Renard, THES-TP 97/02, PM/97-02, hep-ph/9703281, to appear in Phys. Rev. D.
- [25] F.M. Renard and C. Verzegnassi, Phys. Rev. **D52** (1995) 1369, Phys. Rev. **D53** (1996) 1290.
- [26] G.J. Gounaris and C.G. Papadopoulos hep-ph/9612378, to appear in Z.f. Physik.
- [27] A. De Rújula, M.B. Gavela, O. Pène and F.J. Vegas, Nucl. Phys. **B357** (1991) 311; A. Bilal, E. Massó and A. De Rújula Nucl. Phys. **B335** (1991) 549.
- [28] K. Hagiwara and M.L. Stong Z. f. Phys. **C92** (1994) 99; S.Y. Choi and K. Hagiwara, Phys. Lett. **B359** (1995) 369.
- [29] M. Krämer, J. Kühn, M.L. Stong and P.M. Zerwas; Z. f. Phys. **C64** (1994) 21.
- [30] G.J. Gounaris and F.M. Renard, Z. f. Phys. **C69** (1996) 513. G.J. Gounaris, J. Layssac and F.M. Renard, Z. f. Phys. **C65** (1995) 254, Z. f. Phys. **C67** (1995) 542.
- [31] I.F. Ginzburg, G.L. Kotkin, V.G. Serbo and V.I. Telnov, Nucl. Instr. and Meth. **205**, (1983) 47; I.F. Ginzburg, G.L. Kotkin, V.G. Serbo, S.L. Panfil and V.I. Telnov, Nucl. Instr. and Meth. **219**,(1984) 5.
- [32] S.Y. Choi, K. Hagiwara and M.S. Baek, Phys. Rev. **D54** (1996) 6703.
- [33] G.J.Gounaris and G.Tsirigoti, hep-ph/9703446, to appear Phys. Rev. D (Sept. 1997).
- [34] K. Hagiwara, R. Szalapski and D. Zeppenfeld, Phys. Lett. **B318** (1993) 155.





Figure 1: Observability limits in the 6-parameter case; (a) among couplings of set (1), (b) among couplings of set (2); from asymmetries alone (- - -), from asymmetries and integrated observables with a normalization uncertainty of 2% (.....), 20% (———).




Figure 2: Constraints from $e^+e^- \rightarrow b\bar{b}$ observables in the 3-free parameter case, projected on the (f_{qB}, f_{bB}) plane; at LEP2 (without polarization) (*dotted*), at NLC (without polarization) (*solid*), at NLC (with polarization) (*ellipse*).